

Neofitos Stefanides, M.D., P.C.

neostefanidesMD@yahoo.com

44-01 Francis Lewis Blvd L3B
Bayside, NY 11361
Ph: 718.989.8515
Fax: 718.626.0102

1205 Franklin Avenue Suite 150
Garden City, NY 11530
Ph: 718.989.8515
Fax: 718.626.0102

30-74 31st st. 2nd fl.
Astoria, NY 11102
Ph: 718.989.8515
Fax: 718.626.0102

Name: _____ Date: _____
Diagnosis: _____ Date of Surgery: _____

Rehabilitation Protocol: Arthroscopic Meniscus Repair

Phase I (Weeks 0-6)

Weightbearing: As tolerated with crutches

Hinged Knee Brace: worn for 4 weeks post-op

- Locked in full extension for ambulation and sleeping – remove for hygiene and PT

(Weeks 0-2)

- Unlocked for ambulation and removed while sleeping, for hygiene and PT (**Weeks 2-4**)

Range of Motion – AAROM → AROM as tolerated

- **Weeks 0-4:** Full ROM – No weightbearing at flexion angles greater than 90°
- **Weeks 4-6:** Full ROM as tolerated – progress to flexion angles greater than 90°

Therapeutic Exercises

- Quad/Hamstring sets, heel slides, straight leg raises, co-contractions
- Isometric abduction and adduction exercises
- Patellar Mobilizations
- At **4 Weeks:** can begin partial wall-sits – keep knee flexion angle less than 90°

Phase II (Weeks 6-12)

Weightbearing: As tolerated -- discontinue crutch use at 6 weeks

Hinged Knee Brace: Discontinue brace use when patient has achieved full extension with no evidence of extension lag

Range of Motion – Full active ROM

Therapeutic Exercises

- Closed chain extension exercises, Hamstring strengthening
- Lunges – 0-90°, Leg press – 0-90°
- Proprioception exercises
- Begin use of the stationary bicycle

Phase III (Weeks 12-16)

Weightbearing: Full weightbearing with normal gait pattern

Range of Motion – Full/Painless ROM

Neofitos Stefanides, M.D., P.C.

neostefanidesMD@yahoo.com

44-01 Francis Lewis Blvd L3B
Bayside, NY 11361
Ph: 718.989.8515
Fax: 718.626.0102

1205 Franklin Avenue Suite 150
Garden City, NY 11530
Ph: 718.989.8515
Fax: 718.626.0102

30-74 31st st. 2nd fl.
Astoria, NY 11102
Ph: 718.989.8515
Fax: 718.626.0102

Therapeutic Exercises

- Continue with quad and hamstring strengthening
- Focus on single-leg strength
- Begin jogging/running
- Plyometrics and sport-specific drills

Phase IV (Months 4-6)

Gradual return to athletic activity as tolerated

Maintenance program for strength and endurance